

# MACRA: Final Family Medicine Quality Measures

Released October 2016.

MIPS ID Number	NQF/ PQRS	CMS E-Measure ID	Data Submission Method	Measure Type	National Quality Strategy Domain	Measure Title and Description <sup>y</sup>	Measure Steward
<b>7. General Practice/Family Medicine</b>							
* § !	0059 /001	122v5	Claims, Web Interface, Registry, EHR	Intermediate Outcome	Effective Clinical Care	Diabetes: Hemoglobin A1c (HbA1c) Poor Control (>9%) Percentage of patients 18-75 years of age with diabetes who had hemoglobin A1c > 9.0% during the measurement period	National Committee for Quality Assurance

**Notice:** This HHS-approved document has been submitted to the Office of the Federal Register (OFR) for publication and has not yet been placed on public display or published in the Federal Register. This document may vary slightly from the official published document if minor editorial changes have been made during the OFR review process. The document published in the Federal Register is the official HHS-approved document.

MIPS ID Number	NQF/ PQRS	CMS E-Measure ID	Data Submission Method	Measure Type	National Quality Strategy Domain	Measure Title and Description <sup>y</sup>	Measure Steward
<b>7. General Practice/Family Medicine</b>							
§	0081 /005	135v5	Registry, EHR	Process	Effective Clinical Care	Heart Failure (HF): Angiotensin-Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) Therapy for Left Ventricular Systolic Dysfunction (LVSD)  Percentage of patients aged 18 years and older with a diagnosis of heart failure (HF) with a current or prior left ventricular ejection fraction (LVEF) < 40% who were prescribed ACE inhibitor or ARB therapy either within a 12-month period when seen in the outpatient setting OR at each hospital discharge	Physician Consortium for Performance Improvement (PCPI®) Foundation
§	0070 /007	145v5	Registry, EHR	Process	Effective Clinical Care	Coronary Artery Disease (CAD): Beta-Blocker Therapy—Prior Myocardial Infarction (MI) or Left Ventricular Systolic Dysfunction (LVEF <40%)  Percentage of patients aged 18 years and older with a diagnosis of coronary artery disease seen within a 12-month period who also have prior MI OR a current or prior LVEF < 40% who were prescribed beta-blocker therapy	Physician Consortium for Performance Improvement Foundation (PCPI®)
* §	0083 /008	144v5	Registry, EHR	Process	Effective Clinical Care	Heart Failure (HF): Beta-Blocker Therapy for Left Ventricular Systolic Dysfunction (LVSD)  Percentage of patients aged 18 years and older with a diagnosis of heart failure (HF) with a current or prior left ventricular ejection fraction (LVEF) < 40% who were prescribed beta-blocker therapy either within a 12-month period when seen in the outpatient setting OR at each hospital discharge	Physician Consortium for Performance Improvement Foundation (PCPI®)
	105/ 009	128v5	EHR	Process	Effective Clinical Care	Anti-Depressant Medication Management  Percentage of patients 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression, and who remained on antidepressant medication treatment. Two rates are reported a. Percentage of patients who remained on an antidepressant medication for at least 84 days (12 weeks) b. Percentage of patients who remained on an antidepressant medication for at least 180 days (6 months)	National Committee for Quality Assurance
	0326 / 047	N/A	Claims, Registry	Process	Communicati on and Care Coordination	Care Plan  Percentage of patients aged 65 years and older who have an advance care plan or surrogate decision maker documented in the medical record or documentation in the medical record that an advance care plan was discussed but the patient did not wish or was not able to name a surrogate decision maker or provide an advance care plan.	National Committee for Quality Assurance

MIPS ID Number	NQF/ PQRS	CMS E-Measure ID	Data Submission Method	Measure Type	National Quality Strategy Domain	Measure Title and Description <sup>x</sup>	Measure Steward
<b>7. General Practice/Family Medicine</b>							
!	N/A/ 050	N/A	Claims, Registry	Process	Person and Caregiver- Centered Experience and Outcomes	Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older  Percentage of female patients aged 65 years and older with a diagnosis of urinary incontinence with a documented plan of care for urinary incontinence at least once within 12 months	National Committee for Quality Assurance
!!	0069 /065	154v5	Registry, EHR	Process	Efficiency and Cost Reduction	Appropriate Treatment for Children with Upper Respiratory Infection (URI)  Percentage of children 3 months through 18 years of age who were diagnosed with upper respiratory infection (URI) and were not dispensed an antibiotic prescription on or three days after the episode	National Committee for Quality Assurance
* !!	N/A/ 066	146v5	Registry, EHR	Process	Efficiency and Cost Reduction	Appropriate Testing for Children with Pharyngitis  Percentage of children 3-18 years of age who were diagnosed with pharyngitis, ordered an antibiotic and received a group A streptococcus (strep) test for the episode	National Committee for Quality Assurance
!!	0654 /093	N/A	Claims, Registry	Process	Efficiency and Cost Reduction	Acute Otitis Externa (AOE): Systemic Antimicrobial Therapy – Avoidance of Inappropriate Use  Percentage of patients aged 2 years and older with a diagnosis of AOE who were not prescribed systemic antimicrobial therapy	American Academy of Otolaryngolog y-Head and Neck Surgery
!	N/A/ 109	N/A	Claims, Registry	Process	Person and Caregiver- Centered Experience and Outcomes	Osteoarthritis (OA): Function and Pain Assessment  Percentage of patient visits for patients aged 21 years and older with a diagnosis of osteoarthritis (OA) with assessment for function and pain	American Academy of Orthopedic Surgeons
	0041 /110	147v6	Claims, Web Interface, Registry, EHR	Process	Community/ Population Health	Preventive Care and Screening: Influenza Immunization  Percentage of patients aged 6 months and older seen for a visit between October 1 and March 31 who received an influenza immunization OR who reported previous receipt of an influenza immunization	Physician Consortium for Performance Improvement Foundation (PCPI®)
* §	2372 /112	125v5	Claims, Web Interface, Registry, EHR	Process	Effective Clinical Care	Breast Cancer Screening  Percentage of women 50 -74 years of age who had a mammogram to screen for breast cancer	National Committee for Quality Assurance
§	0034 /113	130v5	Claims, Web Interface, Registry, EHR	Process	Effective Clinical Care	Colorectal Cancer Screening  Percentage of patients 50 - 75 years of age who had appropriate screening for colorectal cancer	National Committee for Quality Assurance

MIPS ID Number	NQF/ PQRS	CMS E-Measure ID	Data Submission Method	Measure Type	National Quality Strategy Domain	Measure Title and Description <sup>x</sup>	Measure Steward
<b>7. General Practice/Family Medicine</b>							
§ !!	0058 /116	N/A	Registry	Process	Efficiency and Cost Reduction	Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis: Percentage of adults 18-64 years of age with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription	National Committee for Quality Assurance
§	0055 /117	131v5	Claims, Web Interface, Registry, EHR	Process	Effective Clinical Care	Diabetes: Eye Exam  Percentage of patients 18 - 75 years of age with diabetes who had a retinal or dilated eye exam by an eye care professional during the measurement period or a negative retinal exam (no evidence of retinopathy) in the 12 months prior to the measurement period	National Committee for Quality Assurance
* §	0062 /119	134v4	Registry, EHR	Process	Effective Clinical Care	<b>Diabetes: Medical Attention for Nephropathy:</b> The percentage of patients 18-75 years of age with diabetes who had a nephropathy screening test or evidence of nephropathy during the measurement period  <b>Rationale:</b> CMS is finalizing MIPS #119 for 2017 Performance Period.	National Committee for Quality Assurance
	0421 /128	69v5	Claims, Registry, EHR, Web Interface	Process	Community/ Population Health	Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan  Percentage of patients aged 18 years and older with a BMI documented during the current encounter or during the previous six months AND with a BMI outside of normal parameters, a follow-up plan is documented during the encounter or during the previous six months of the current encounter	Centers for Medicare & Medicaid Services
	0419 /130	68v6	Claims, Registry, EHR,	Process	Patient Safety	Documentation of Current Medications in the Medical Record  Percentage of visits for patients aged 18 years and older for which the eligible professional attests to documenting a list of current medications using all immediate resources available on the date of the encounter. This list must include ALL known prescriptions, over-the-counters, herbals, and vitamin/mineral/dietary (nutritional) supplements AND must contain the medications' name, dosage, frequency and route of administration.	Centers for Medicare & Medicaid Services
*	0418 /134	2v6	Claims, Web Interface, Registry, EHR	Process	Community/ Population Health	Preventive Care and Screening: Screening for Depression and Follow-Up Plan  Percentage of patients aged 12 years and older screened for depression on the date of the encounter using an age appropriate standardized depression screening tool AND if positive, a follow-up plan is documented on the date of the positive screen	Centers for Medicare & Medicaid Services

MIPS ID Number	NQF/ PQRS	CMS E-Measure ID	Data Submission Method	Measure Type	National Quality Strategy Domain	Measure Title and Description <sup>x</sup>	Measure Steward
<b>7. General Practice/Family Medicine</b>							
!	0101 /154	N/A	Claims, Registry	Process	Patient Safety	Falls: Risk Assessment  Percentage of patients aged 65 years and older with a history of falls who had a risk assessment for falls completed within 12 months	National Committee for Quality Assurance
!	0101 /155	N/A	Claims, Registry	Process	Communication and Care Coordination	Falls: Plan of Care  Percentage of patients aged 65 years and older with a history of falls who had a plan of care for falls documented within 12 months	National Committee for Quality Assurance
* §	0056 /163	123v5	EHR	Process	Effective Clinical Care	Comprehensive Diabetes Care: Foot Exam  The percentage of patients 18-75 years of age with diabetes (type 1 and type 2) who received a foot exam (visual inspection and sensory exam with mono filament and a pulse exam) during the measurement year	National Committee for Quality Assurance
!	NA/ 181	N/A	Claims, Registry	Process	Patient Safety	Elder Maltreatment Screen and Follow-Up Plan  Percentage of patients aged 65 years and older with a documented elder maltreatment screen using an Elder Maltreatment Screening Tool on the date of encounter AND a documented follow-up plan on the date of the positive screen	Centers for Medicare & Medicaid Services
* §	0068 /204	164v5	Claims, Web Interface, Registry, EHR	Process	Effective Clinical Care	Ischemic Vascular Disease (IVD): Use of Aspirin or Another Antiplatelet  Percentage of patients 18 years of age and older who were diagnosed with acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) in the 12 months prior to the measurement period, or who had an active diagnosis of ischemic vascular disease (IVD) during the measurement period, and who had documentation of use of aspirin or another antiplatelet during the measurement period	National Committee for Quality Assurance
	0028 /226	138v5	Claims, Registry, EHR, Web Interface	Process	Community/Population Health	Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention  Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user.	Physician Consortium for Performance Improvement Foundation (PCPI®)
	0018 /236	165v5	Claims, Registry, EHR, Web Interface	Intermediate Outcome	Effective Clinical Care	Controlling High Blood Pressure  Percentage of patients 18-85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (<140/90 mmHg) during the measurement period	National Committee for Quality Assurance

MIPS ID Number	NQF/ PQRS	CMS E-Measure ID	Data Submission Method	Measure Type	National Quality Strategy Domain	Measure Title and Description <sup>x</sup>	Measure Steward
<b>7. General Practice/Family Medicine</b>							
* §	0032 /309	124v5	EHR	Process	Effective Clinical Care	Cervical Cancer Screening  Percentage of women 21–64 years of age who were screened for cervical cancer using either of the following criteria: <ul style="list-style-type: none"> <li>• Women age 21–64 who had cervical cytology performed every 3 years</li> <li>• Women age 30–64 who had cervical cytology/human papillomavirus (HPV) co-testing performed every 5 years</li> </ul>	National Committee for Quality Assurance
§ !!	0052 /312	166v6	EHR	Process	Efficiency and Cost Reduction	Use of Imaging Studies for Low Back Pain  Percentage of patients 18-50 years of age with a diagnosis of low back pain who did not have an imaging study (plain X-ray, MRI, CT scan) within 28 days of the diagnosis	National Committee for Quality Assurance
*	N/A/ 317	22v5	Claims, Registry, EHR	Process	Community/P opulation Health	Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented  Percentage of patients aged 18 years and older seen during the reporting period who were screened for high blood pressure AND a recommended follow-up plan is documented based on the current blood pressure (BP) reading as indicated.	Centers for Medicare & Medicaid Services
§ !	0005 & 0006 /321	N/A	CMS- approved Survey Vendor	Patient Engagemen t/Experienc e	Person and Caregiver- Centered Experience and Outcomes	CAHPS for MIPS Clinician/Group Survey:  <u>Summary Survey Measures may include:</u> <ul style="list-style-type: none"> <li>• Getting Timely Care, Appointments, and Information;</li> <li>• How well Providers Communicate;</li> <li>• Patient’s Rating of Provider;</li> <li>• Access to Specialists;</li> <li>• Health Promotion and Education;</li> <li>• Shared Decision-Making;</li> <li>• Health Status and Functional Status;</li> <li>• Courteous and Helpful Office Staff;</li> <li>• Care Coordination;</li> <li>• Between Visit Communication;</li> <li>• Helping You to Take Medication as Directed; and</li> <li>• Stewardship of Patient Resources.</li> </ul>	
§	1525 /326	N/A	Claims, Registry	Process	Effective Clinical Care	Atrial Fibrillation and Atrial Flutter: Chronic Anticoagulation Therapy  Percentage of patients aged 18 years and older with a diagnosis of nonvalvular atrial fibrillation (AF) or atrial flutter whose assessment of the specified thromboembolic risk factors indicate one or more high-risk factors or more than one moderate risk factor, as determined by CHADS2 risk stratification, who are prescribed warfarin OR another oral anticoagulant drug that is FDA approved for the prevention of thromboembolism	American College of Cardiology

MIPS ID Number	NQF/ PQRS	CMS E-Measure ID	Data Submission Method	Measure Type	National Quality Strategy Domain	Measure Title and Description <sup>x</sup>	Measure Steward
<b>7. General Practice/Family Medicine</b>							
!!	N/A/ 331	N/A	Registry	Process	Efficiency and Cost Reduction	Adult Sinusitis: Antibiotic Prescribed for Acute Sinusitis (Overuse)  Percentage of patients, aged 18 years and older, with a diagnosis of acute sinusitis who were prescribed an antibiotic within 10 days after onset of symptoms	American Academy of Otolaryngology-Head and Neck Surgery
!!	N/A/ 332	N/A	Registry	Process	Efficiency and Cost Reduction	Adult Sinusitis: Appropriate Choice of Antibiotic: Amoxicillin With or Without Clavulanate Prescribed for Patients with Acute Bacterial Sinusitis (Appropriate Use)  Percentage of patients aged 18 years and older with a diagnosis of acute bacterial sinusitis that were prescribed amoxicillin, with or without clavulanate, as a first line antibiotic at the time of diagnosis	American Academy of Otolaryngology-Head and Neck Surgery
!!	N/A/ 333	N/A	Registry	Efficiency	Efficiency and Cost Reduction	Adult Sinusitis: Computerized Tomography (CT) for Acute Sinusitis (Overuse)  Percentage of patients aged 18 years and older with a diagnosis of acute sinusitis who had a computerized tomography (CT) scan of the paranasal sinuses ordered at the time of diagnosis or received within 28 days after date of diagnosis	American Academy of Otolaryngology-Head and Neck Surgery
!!	N/A/ 334	N/A	Registry	Efficiency	Efficiency and Cost Reduction	Adult Sinusitis: More than One Computerized Tomography (CT) Scan Within 90 Days for Chronic Sinusitis (Overuse)  Percentage of patients aged 18 years and older with a diagnosis of chronic sinusitis who had more than one CT scan of the paranasal sinuses ordered or received within 90 days after the date of diagnosis	American Academy of Otolaryngology-Head and Neck Surgery
	N/A/ 337	N/A	Registry	Process	Effective Clinical Care	Tuberculosis (TB) Prevention for Psoriasis, Psoriatic Arthritis and Rheumatoid Arthritis Patients on a Biological Immune Response Modifier  Percentage of patients whose providers are ensuring active tuberculosis prevention either through yearly negative standard tuberculosis screening tests or are reviewing the patient's history to determine if they have had appropriate management for a recent or prior positive test	American Academy of Dermatology
* § !	2082 /338	N/A	Registry	Outcome	Effective Clinical Care	HIV Viral Load Suppression  The percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year	Health Resources and Services Administration

MIPS ID Number	NQF/ PQRS	CMS E-Measure ID	Data Submission Method	Measure Type	National Quality Strategy Domain	Measure Title and Description <sup>x</sup>	Measure Steward
<b>7. General Practice/Family Medicine</b>							
!	N/A/ 342	N/A	Registry	Outcome	Person and Caregiver- Centered Experience and Outcomes	Pain Brought Under Control Within 48 Hours  Patients aged 18 and older who report being uncomfortable because of pain at the initial assessment (after admission to palliative care services) who report pain was brought to a comfortable level within 48 hours	National Hospice and Palliative Care Organization
* § !	0710 /370	159v5	Web Interface, Registry, EHR	Outcome	Effective Clinical Care	Depression Remission at Twelve Months:  Patients age 18 and older with major depression or dysthymia and an initial Patient Health Questionnaire (PHQ-9) score greater than nine who demonstrate remission at twelve months (+/- 30 days after an index visit) defined as a PHQ-9 score less than five. This measure applies to both patients with newly diagnosed and existing depression whose current PHQ-9 score indicates a need for treatment.	Minnesota Community Measurement
	N/A/ 387	N/A	Registry	Process	Effective Clinical Care	Annual Hepatitis C Virus (HCV) Screening for Patients who are Active Injection Drug Users  Percentage of patients regardless of age who are active injection drug users who received screening for HCV infection within the 12 month reporting period	Physician Consortium for Performance Improvement Foundation (PCPI®)
	1407 /394	N/A	Registry	Process	Community/ Population Health	Immunizations for Adolescents  The percentage of adolescents 13 years of age who had the recommended immunizations by their 13th birthday	National Committee for Quality Assurance
!	N/A/ 398	N/A	Registry	Outcome	Effective Clinical Care	Optimal Asthma Control  Composite measure of the percentage of pediatric and adult patients whose asthma is well-controlled as demonstrated by one of three age appropriate patient reported outcome tools	Minnesota Community Measurement
§	N/A/ 400	N/A	Registry	Process	Effective Clinical Care	One-Time Screening for Hepatitis C Virus (HCV) for Patients at Risk  Percentage of patients aged 18 years and older with one or more of the following: a history of injection drug use, receipt of a blood transfusion prior to 1992, receiving maintenance hemodialysis OR birthdate in the years 1945-1965 who received one-time screening for hepatitis C virus (HCV) infection	Physician Consortium for Performance Improvement Foundation (PCPI®)



MIPS ID Number	NQF/ PQRS	CMS E-Measure ID	Data Submission Method	Measure Type	National Quality Strategy Domain	Measure Title and Description <sup>x</sup>	Measure Steward
<b>7. General Practice/Family Medicine</b>							
§	N/A/ 401	N/A	Registry	Process	Effective Clinical Care	Hepatitis C: Screening for Hepatocellular Carcinoma (HCC) in Patients with Cirrhosis  Percentage of patients aged 18 years and older with a diagnosis of chronic hepatitis C cirrhosis who underwent imaging with either ultrasound, contrast enhanced CT or MRI for hepatocellular carcinoma (HCC) at least once within the 12 month reporting period	American Gastroenterological Association/ American Society for Gastrointestinal Endoscopy/ American College of Gastroenterology
	NA/ 402	NA	Registry	Process	Community/ Population Health	Tobacco Use and Help with Quitting Among Adolescents  The percentage of adolescents 12 to 20 years of age with a primary care visit during the measurement year for whom tobacco use status was documented and received help with quitting if identified as a tobacco user	National Committee for Quality Assurance
	N/A/ 408	N/A	Registry	Process	Effective Clinical Care	Opioid Therapy Follow-up Evaluation  All patients 18 and older prescribed opiates for longer than six weeks duration who had a follow-up evaluation conducted at least every three months during Opioid Therapy documented in the medical record	American Academy of Neurology
	N/A/ 412	N/A	Registry	Process	Effective Clinical Care	Documentation of Signed Opioid Treatment Agreement  All patients 18 and older prescribed opiates for longer than six weeks duration who signed an opioid treatment agreement at least once during Opioid Therapy documented in the medical record	American Academy of Neurology
	N/A/ 414	N/A	Registry	Process	Effective Clinical Care	Evaluation or Interview for Risk of Opioid Misuse  All patients 18 and older prescribed opiates for longer than six weeks duration evaluated for risk of opioid misuse using a brief validated instrument (e.g. Opioid Risk Tool, SOAAP-R) or patient interview documented at least once during Opioid Therapy in the medical record	American Academy of Neurology
	0053 /418	N/A	Claims, Registry	Process	Effective Clinical Care	Osteoporosis Management in Women Who Had a Fracture  The percentage of women age 50-85 who suffered a fracture and who either had a bone mineral density test or received a prescription for a drug to treat osteoporosis in the six months after the fracture	National Committee for Quality Assurance

MIPS ID Number	NQF/ PQRS	CMS E-Measure ID	Data Submission Method	Measure Type	National Quality Strategy Domain	Measure Title and Description <sup>x</sup>	Measure Steward
<b>7. General Practice/Family Medicine</b>							
	2152 /431	NA	Registry	Process	Community/ Population Health	Preventive Care and Screening: Unhealthy Alcohol Use: Screening & Brief Counseling  Percentage of patients aged 18 years and older who were screened for unhealthy alcohol use using a systematic screening method at least once within the last 24 months AND who received brief counseling if identified as an unhealthy alcohol user.	Physician Consortium for Performance Improvement Foundation (PCPI®)
	N/A/ 438	N/A	Web Interface, Registry	Process	Effective Clinical Care	Statin Therapy for the Prevention and Treatment of Cardiovascular Disease  Percentage of the following patients—all considered at high risk of cardiovascular events—who were prescribed or were on statin therapy during the measurement period: • Adults aged ≥ 21 years who were previously diagnosed with or currently have an active diagnosis of clinical atherosclerotic cardiovascular disease (ASCVD); OR • Adults aged ≥21 years with a fasting or direct low-density lipoprotein cholesterol (LDL-C) level ≥ 190 mg/dL; OR • Adults aged 40-75 years with a diagnosis of diabetes with a fasting or direct LDL-C level of 70-189 mg/dL	Centers for Medicare & Medicaid Services
<b>+ §</b>	0071 /442	N/A	Registry	Process	Effective Clinical Care	Persistent Beta Blocker Treatment After a Heart Attack  The percentage of patients 18 years of age and older during the measurement year who were hospitalized and discharged from July 1 of the year prior to the measurement year to June 30 of the measurement year with a diagnosis of acute myocardial infarction (AMI) and who received were prescribed persistent beta-blocker treatment for six months after discharge	National Committee for Quality Assurance
<b>+ § !!</b>	N/A/ 443	N/A	Registry	Process	Patient Safety	Non-Recommended Cervical Cancer Screening in Adolescent Females  The percentage of adolescent females 16–20 years of age screened unnecessarily for cervical cancer	National Committee for Quality Assurance
<b>+ § !</b>	1799 /444	NA	Registry	Process	Efficiency and Cost Reduction	Medication Management for People with Asthma (MMA):  The percentage of patients 5-64 years of age during the measurement year who were identified as having persistent asthma and were dispensed appropriate medications that they remained on for at least 75% of their treatment period.	National Committee for Quality Assurance

MIPS ID Number	NQF/ PQRS	CMS E-Measure ID	Data Submission Method	Measure Type	National Quality Strategy Domain	Measure Title and Description <sup>x</sup>	Measure Steward
<b>7. General Practice/Family Medicine</b>							
<p><b>Comment:</b> CMS received specific comments to add several individual measures and cross-cutting measures to the measure set because the commenters believed the additional measures were appropriate for providers within the general practice and family medicine specialties. Commenters specifically asked that measures #007, #008, #046, #047, #110, #119, #163, #204, #226, #236, #309, #321, #370, #442, #443 and #444 be added to the measure set.</p> <p><b>Response:</b> Upon further review of the recommendations provided by commenters, CMS has revised the measure set from the proposed set by adding these relevant measures to the measures set (#007, #008, #047, #110, #119, #163, #204, #226, #309, #321, #370, #442, #443, and #444). CMS did not include measure #46 in the General Practice measure set because we are including measure #130, a cross-cutting measure, which is closely related to this measure, to the set. In addition, CMS has added previously identified cross-cutting measures that are relevant for the specialty set (#047, #128, #130, #226, #317, #402, and #431). CMS believes the finalized specialty set reflects the relevant measures appropriate for the family medicine/general practice specialty.</p> <p><b>Final Decision:</b> CMS is finalizing the general practice and family medicine specialty measure set as indicated in the table above.</p>							